

CURRENT RESEARCH
IN
MUSCULAR DYSTROPHY, JAPAN
(CLINICAL RESEARCHES)

The Proceedings of the
Annual Meeting of Muscular
Dystrophy Research Group,
1980, Tokyo

PREFACE

Following the last Proceedings of the Annual Meeting of Muscular Dystrophy Research Group, 1978, 1979, "Current Research in Muscular Dystrophy, 1980" is published.

This volume contains not only the abstracts of the papers presented at the Annual Meeting of the Clinical Research Group of Muscular Dystrophy in 1980, but also some of the research works performed during past three years.

Meanwhile, a real progress has been made in the fields of the genetic, clinical, morphological and metabolic aspects of the disease and this may lead to the understanding of the fundamental defects. Still, we have to emphasize that these and other discoveries, while they are significantly advanced and certainly very encouraging, are by no means final.

Lastly, I as the chairman of this Research Group, would like to express my sincere appreciation to the Ministry of Health and Welfare, National Center for Nervous, Mental and Muscular Disorders, and Muscular Dystrophy Association of Japan for persistent financial support to this Research Group for three years.

My thanks are also due to Dr. H. Sugita, the co-editor, who has made it possible to publish this report.

Kazuo Miyoshi

March, 1981

CONTENTS

PREFACE

[CHAPTER ONE] Experimental Muscular Dystrophy in Chicken

- 1) Early Morphological Changes in Dystrophic Chicken Muscles:
Yoshigoro Kuroiwa 10
- 2) Morphometric Investigation of Developing Muscles of Dystrophic
Chicken with Special Reference to the Posterior Latissimus Dorsi Muscle:
Haruomi Nakamura 12
- 3) Intracytoplasmic Vacuolation of α W Fibers in Chicken Muscular Dystrophy
– Probable Early Pathological Event Initiates Massive Fiber Necrosis –:
Ikuya Nonaka 14
- 4) A Radioautographic Study of the Activated Satellite Cell in Dystrophic
Chicken Muscle: Kimio Oguchi 16
- 5) Trophic Effects of Sympathetic Ganglia on Normal and Dystrophic Chick
Skeletal Muscles in Tissue Culture (2) – Acetylcholinesterase and
Catecholamine Staining –: Hiroshi Tsukagoshi 18
- 6) The Movement of “Filopodia” to Normal and Dystrophic Chick Muscles:
Takao Nakanishi 20
- 7) Study on Neutral Lipid in the Muscle of Dystrophic Chicken:
Masao Kinoshita 22
- 8) Gangliosides in Dystrophic Chicken Muscle: Tadashi Miyatake 24

[CHAPTER TWO] Genetic and Clinical Studies of Progressive Muscular Dystrophy and Related Disorders

- 1) Genetic Analysis of Amyotrophic Diseases – Duchenne Type Muscular
Dystrophy –: Toshiyuki Furusho 26
- 2) Facioscapulohumeral Syndrome: The Non-Hereditary Variety:
Tetsuo Furukawa 28
- 3) Familial Oculocranoskeletal Neuromuscular Disease with Abnormal
Muscle Mitochondria – A Follow Up Study: Shukuro Araki 30
- 4) Computerized Tomographic Findings of the Central Nervous System in
Myotonic Dystrophy: Hiroo Imura 34
- 5) An Unusual Case with Facio-Scapulo-Humeral Muscular atrophy:
Keishi Ueda 36
- 6) Centronuclear Myopathy – A Distal from with A Distal from with Juvenile
Onset and No Ocular Involvement: Keiichi Takahashi 38
- 7) A Case of Atypical Myopathy with Contractures of Joints and Cardiac
Involvement: Hiroshi Tsukagoshi 40

[CHAPTER THREE] Anatomical and Pathological Studies of Progressive Muscular Dystrophy and Related Disorders

1) Muscle Pathology of Distal Myopathy: Yasuo Toyokura	42
2) Distal Myopathy and Lysosomes: Nobuyoshi Fukuhara	44
3) Tubulomembranous Myopathy: Nobuyoshi Fukuhara	46
4) Debrancher Deficiency Myopathy: Three Adults Including Two Siblings: Mitsuhiro Osame	48
5) Transplantation of Human Dystrophic Muscle to Nude Mice: A Morphologic Study: Yoshihiro Wakayama	50
6) A Morphological Study of Experimental Mitochondrial Myopathy with 2, 4-Dinitrophenol: Shukuro Araki	52
7) Myopathy with Mitochondrial Inclusion Bodies – Clinical and Electron- microscopic Studies –: Takeshi Sato	54
8) Study on Disuse Atrophy: Masao Kinoshita	58
9) Histochemical Study on Lipid Metabolism in Motor Neurone Disease: Toshiji Mozai	60
10) Histological Studies on Muscle in a Case with Schwartz-Jampel Syndrome—: Toshiji Mozai	62

[CHAPTER FOUR] Physiological Studies of Progressive Muscular Dystrophy and Related Disorders

1) Single Fiber EMG of Progressive Muscular Dystrophies: Hiroschi Nishitani	64
2) Experimental Myotonia in Human External Intercostal Muscles: Teruyuki Kurihara	66
3) Contractile and Chemosensitive Properties of Muscle Treated with Calcium Ionophore (A23187): Masaharu Takamori	68

[CHAPTER FIVE] Biochemical Studies of Progressive Muscular Dystrophy and Related Disorders

a) Reds Cells	
1) Phosphatidyl Inositol Metabolism in Duchenne Erythrocyte Membranes: Keiichi Takahashi	70
2) Concanavalin A Binding on the Erythrocytes in Duchenne Muscular Dystrophy: Masanori Uono	72
3) Calcium Flux of Erythrocytes in Duchenne Muscular Dystrophy: Shin'ichi Shoji	74
4) Ca and ATP Contents of Erythrocytes in Duchenne Muscular Dystrophy: Mizuko Yoshida	76
5) ATPase and Adenylate Kinase of Erythrocyte Membranes in Human Myotonic Dystrophy: Shiro Mawatari	78

6) Erythrocyte Acetylcholinesterase in Myotonic Dystrophy: Yoshigoro Kuroiwa	80
b) Metabolism and Enzymes	
1) Congenital Myopathy and Abnormal Tryptophan Metabolism: Masatoshi Yagita	82
2) Some Properties of Creatinase in Human Skeletal Muscle: Yasuo Natori	84
3) Abnormal Metabolism of Creatine in Skeletal Muscles of Muscular Dystrophies: Kazuo Miyoshi	86
4) Enzymatic Studies on the Mechanism of Creatinuria in Rats with Experimental Thyrotoxic Myopathy: Keishi Ueda	88
5) Exertional Rhabdomyolysis and Myoglobinuria: A Case Report and Metabolic Study: Tadashi Miyatake	92
6) Turnover of Human Myoglobin as Studied by Radioimmunoassay: Hisaoami Kawai	94
7) GOT Isozymes in Muscles of Patients with Ophthalmoplegia Plus Syndrome and Experimental Mitochondrial Myopathy: Seiichiro Tarui	96
c) Contractile Proteins and Proteases	
1) Analysis of the Contractile Protein of a Single Human Muscle Fiber: Akio Takagi	100
2) Studies on the Structural Proteins in Experimental Serotonin Myopathy: Mitsuo Iida	102
3) Calcium-Dependent Degradation in Muscle Protein – Effect of Ca Ionophore (A23187) and a CANP Inhibitor (E-64-c) –: Hideo Sugita	104
4) Degradation of Myofibrillar Proteins by Cathepsin H: Yukihiro Sanada	106
d) Others	
1) Immunoglobulin and the Immune Complex in Fukuyama Type Congenital Muscular Dystrophy: Kiyotaro Kondo	108
2) Isolation of Nerve Growth Factor from Human Placental Tissue: Kyozo Hayashi	110
3) Level of Nerve Growth Factor-Like Immunoreactivity in Human Serum: Kyozo Hayashi	112

THE LIST OF THE MEMBERS OF THE CLINICAL RESEARCH GROUP ON MUSCULAR DYSTROPHY IN JAPAN	115
--	------------

The Proceedings of the
Annual Meeting of Muscular
Dystrophy Research Group,
1980, Tokyo

EARLY MORPHOLOGICAL CHANGES IN DYSTROPHIC CHICKEN MUSCLES

Yoshigoro Kuroiwa, Shinji Ishimoto, Ikuo Goto and Michiya Ohta*

Department of Neurology and Department of Neuropathology*, Neurological Institute, Faculty of Medicine, Kyushu University, Fukuoka

We previously reported¹⁾ that early morphological changes appeared in PLD muscles at 1 week after hatching and these changes became more prominent at 3 weeks after hatching. This time, we studied the caveolae on the plasma membrane surface of the superficial pectoral muscles at various stages after hatching in control and dystrophic chickens.

Materials and methods

We used the superficial pectoral muscles, white fibers, of control (line 412) and dystrophic chickens (line 413) at 1 day, 1 week and 3 weeks after hatching. For the freeze-fracture EM study, the muscle specimens were routinely fixed in 2.5% glutaraldehyde and cut into 1-mm cubes following fixation. Then, the blocks were immersed in 25% glycerol. These specimens were frozen by immersion in liquid nitrogen. They were fractured and shadowed with platinum and carbon in a freeze-fracture device operated at -115°C and beneath 1×10^{-6} Torr. Replicas were cleaned in bleach and washed with distilled

water and examined with an electron microscope. 200-micron square areas representing a total sampling area were counted in order to determine average caveolar density.

Results (Fig. 1)

At 1 day and 1 week after hatching, there was a difference in caveolar density between control and dystrophic muscles. The distribution, shape and size of caveolae were more irregular in dystrophic chickens (Fig. 2). There were no remarkable differences in caveolar density between 1 day and 1 week after hatching. However, the caveolar distribution showed more random arrangement in dystrophic muscles. At 3 weeks after hatching, there was an increase of caveolar density in dystrophic muscles as compared with control muscles and the caveolar distribution was more irregular and random in dystrophic muscles. In control muscles, the caveolae were arranged in a regular pattern. There were no remarkable changes in caveolar density of control muscles at 1 day, 1 week or 3 weeks after hatching.

Discussion

The caveolae are considered to be either pinocytotic vesicles or T-system openings. There were no obvious changes in dystrophic




Fig. 1 CHANGES OF CAVEOLAR DENSITY (total sampling areas: $200 \mu\text{m}^2$)

muscles at 1 day after hatching in our previous study, while the changes of caveolar density, distribution and size were already seen at 1 day after hatching in dystrophic muscles. This study of caveolae is considered to be more useful for detecting the early morphological changes of the dystrophic process.

References

- 1) Kuroiwa, Y., Ishimoto, S., Goto, I. and Ohta, M.: Early morphological changes in dystrophic chicken muscle. Proceedings of Ann. Meet. of Muscular Dystrophy Res. Group., p. 12, 1978 and 1979.




Fig. 2

- (C) P face of muscle plasmalemma from a 1 day old normal chicken. The caveolar distribution has a regular pattern. $\times 21,000$
- (D) E face of muscle plasmalemma from a 1 day old dystrophic chicken. The caveolar distribution is more random and the distribution and size are more variable than in control chickens. $\times 40,000$