

CURRENT RESEARCH
IN
MUSCULAR DYSTROPHY, JAPAN
(BIOMEDICAL RESEARCHES)

Vol. 3

The Proceedings of the
Annual Meeting of Muscular
Dystrophy Research Group,
1981, Tokyo

CONTENTS

Structural Changes in Striated Muscles of Dystrophic Hamster	1
Cell Adhesion Molecule of Cleavage Stage Mouse Embryos—Functions in “Compaction”	3
Phage Particle-Mediated Gene Transfer into Cultured Mammalian Cells	5
A Study of Dystrophic Muscle Maintained <i>in vitro</i>	7
Mitotic Activities of Mononucleate Cells Cultured <i>in vitro</i> Originated from Regenerating Muscles of Normal (+/+) and Dystrophic (dy/dy) Mice (II)	9
Requirement of Transferrin for Myogenic Cell Growth <i>in vitro</i>	11
Effects of Innervation on Muscle Regeneration	13
Postnatal Alterations in the Number of Muscle Fibers in Normal and Dystrophic Chickens	15
Arrested Growth of Muscle Fibers in Muscular Dystrophic Mice	17
Comparison of the Alteration Patterns of Several Biochemical Markers in Mice with Some Neuromuscular Mutations	19
Effect of Brain Extract on Development of Tetrodotoxin-Sensitive Sodium Channel in L6 Myotubes	21
Developmental Studies of the Membrane Properties of Striated Muscle Fibers—Changes in the Properties of Ca Channels during Differentiation of Excitable Cells	23
Sensitivity to Ca ²⁺ , Sr ²⁺ , and Ba ²⁺ Ions of the Contractile System of Slow and Fast Fibers of Amphibian Muscles	25
Antibodies to Acetylcholine Receptor in Ocular Myasthenia Gravis	27
Neuronal Factor Stabilizing Acetylcholine Receptors on Cultured Muscle Cells	29
Synthesis and Differentiation of Acetylcholine Receptors in Cultured Skeletal Muscle Cells	31
Studies on Acetylcholine Receptor of Dystrophic Chick Muscle in Culture	33
Freeze-Fracture Analysis of the Caveolae in Chick Skeletal Muscles	35
Freeze-Etch Replica Study of the Cytoskeleton in Skeletal Muscle Fibers	37
Different Rate of Change in Isoforms of Myosin Heavy and Light Chains during Development of Skeletal Muscle	39
Number of Antitroponin I Striations along the Thin Filament of Adult and Fetal Skeletal Muscles of Rabbit	41
Isoform Variants of Myofibrillar Proteins during Muscle Regeneration after Cold Injury	43
Myosin Isozymes in Embryonic and Adult Chicken Skeletal Muscle	45
Native High Molecular Weight Protein from Chicken Breast Muscle	47
A Protein in Chicken Breast Muscle Which Inhibits the ADP-Ribosylation of EF2 by Diphtheria Toxin <i>in vitro</i>	49
A Trophic Interaction in Denervated Muscles from the Standpoint of Recovery of Ca-Uptake Ability of FSR	51
Changes in Protease Activities during Development of Chicken Muscles	53
Endogenous Thiol Proteinase Inhibitor and Its Changes of Activity in Skeletal Muscle of Dystrophic Hamsters	55
The Effects of Low Molecular Enzyme Inhibitors on Enzyme Networks in Various Organs of Dystrophic Mice	57
The Role of Protease in the Development and Differentiation of Muscles	59
The Effect of Protease Inhibitor on the Calcium-Induced Degeneration of Myofilament <i>in vitro</i> and <i>in vivo</i>	61

STRUCTURAL CHANGES IN STRIATED MUSCLES OF DYSTROPHIC HAMSTER

Yoshiaki Nonomura

Department of Pharmacology, Faculty of Medicine, University of Tokyo, Tokyo 113

Since the first introduction of a strain of myopathic hamster by Homburger's group¹⁾ in 1962, attention has been given to the myopathy of this animal, especially its cardiomyopathy. This strain has an autosomal recessively inherited polymyopathy, but has no other defective organs or tissues except for these striated muscles.

The present work aimed at checking the degree of damage in various skeletal muscles at different ages. BIO 15.6 male hamsters were used for the studies. After being sacrificed by a blow on the head and bleeding, the animal was fixed on a board and the various muscles were carefully removed including both terminal tendons. The muscles were fixed *in situ* length on dental wax and 4% glutaraldehyde buffered with cacodylate was poured on the materials. They were then teased into small bundles of fibers by forceps 30 min later and transferred to a small bottle containing a glutaraldehyde fixative. The usual methods were then followed of a second fixation of osmium tetroxide, staining *en bloc*, dehydration with alcohol and acetone, and embedding in epoxy resin. A relatively thick section with toluidine blue staining was prepared for light microscopy and a thin section with lead staining for electron microscopy. The cross section view for light microscopy was mainly used to determine the grade of muscle damage and this was done by the criteria according to Bajusz *et al.*²⁾, slightly modified as follows: ±, less than 4 cells with central nuclei (CN) among 100 cross sectioned cells; +, less than 8 (CN) cells in 100; ++, more than 8 (CN) cells in 100; +++, abundance of CN cells and other abnormalities, + + + +, severe abnormalities with necrotic damage.

As shown in Table 1, 12 striated muscles of different ages were tested with fixed materials. All had some degree of abnormalities, particularly severe ones in the shoulder muscle. When the central nuclei were seen in a longitudinal section, they were arranged in a long row. In this same section an abnormal arrangement of nuclei was recognized in a long row although they were not localized at the center but at the periphery (Fig. 1). Other nucleic abnormalities such as perinuclear cavities or pores were sometimes present. Myofibrils usually looked generally normal even in a grade +++ muscle, except for the nuclear anomaly. When the muscle was observed in detail under high magnification, both free polysomes and membrane-bound ribosomes were distributed around the nucleus and submembraneous region, which could not be seen in a normal adult muscle. In a grade + + + + muscle some vacuoles and disarrangement of myofibrils appeared here and there together with abundant CN.

TABLE 1. The Degree of Muscle Damage

	77 days	97 days	110 days	130 days
Arm muscles M. extensor digitrum communis				±
M. extensor carpi radialis brevis				++
Shoulder muscles M. omotransversarius		+++*	++++*	
M. omobrachialis				+++
Back muscle M. lumbodorsal aponeurosis				++++*
Abdominal muscle M. external abdominal oblique			+	
Esophagus M. esophagus		+	+	
Diaphragm M. diaphragma		+++	+++*	
Leg muscles M. samitendinosus		++	+	
M. soleus	++			
M. extensor digitrum longus	+++			
M. sartorium	++		++++*	

* Appearance of necrotic cell.

